

#555

Earth Sc

NIMBUS 5

72-097A-04E

BRIGHTNESS TEMP AND SEA ICE CONCENTRATIONS

ESAD 00218

dec -

*xferred to
NSIDC*

*data not
needed*

Table of Contents

1. Introduction
2. Errata/Change Log
3. LINKS TO RELEVANT INFORMATION IN THE ONLINE NSSDC
INFORMATION SYSTEM
4. Catalog Materials
 - a. Associated Documents
 - b. Core Catalog Materials

1. INTRODUCTION:

The documentation for this data set was originally on paper, kept in NSSDC's Data Set Catalogs (DSCs). The paper documentation in the Data Set Catalogs have been made into digital images, and then collected into a single PDF file for each Data Set Catalog. The inventory information in these DSCs is current as of July 1, 2004. This inventory information is now no longer maintained in the DSCs, but is now managed in the inventory part of the NSSDC information system. The information existing in the DSCs is now not needed for locating the data files, but we did not remove that inventory information.

The offline tape datasets have now been migrated from the original magnetic tape to Archival Information Packages (AIP's).

A prior restoration may have been done on data sets, if a requestor of this data set has questions; they should send an inquiry to the request office to see if additional information exists.

2. ERRATA/CHANGE LOG:

NOTE: Changes are made in a text box, and will show up that way when displayed on screen with a PDF reader.

When printing, special settings may be required to make the text box appear on the printed output.

Version	Date	Person	Page	Description of Change
01				
02				

3 LINKS TO RELEVANT INFORMATION IN THE ONLINE NSSDC
INFORMATION SYSTEM:

<http://nssdc.gsfc.nasa.gov/nmc/>

[NOTE: This link will take you to the main page of the NSSDC Master Catalog. There you will be able to perform searches to find additional information]

4. CATALOG MATERIALS:

- a. Associated Documents To find associated documents you will need to know the document ID number and then click here.
<http://nssdcftp.gsfc.nasa.gov/miscellaneous/documents/>

- b. Core Catalog Materials

REQ. AGENT
DEW

RAND NO.
V0159

ACQ. AGENT
CYN

NIMBUS 5

BRIGHTNESS TEMP AND SEA ICE CONCENTRATIONS

72-097A-04E ESAD-00218

This data set catalog consists of 5 tapes. The tapes are 6250 BPI, 9 track, EBCDIC, and multi-filed. The tapes were created on an IBM computer. The time spans and D and C numbers are as follows:

D#	C#	FILES	TIME SPAN
D-47329	C-22682	140	01/01/73 - 12/31/76
D-79706	C-29056	270	12/19/72 - 12/31/75 Arctic Brightness
D-79707	C-29057	341	12/19/72 - 08/15/76 Antarctic Brightness
D-79708	C-29058	136	12/19/72 - 08/15/76 Arctic Brightness Tem
D-79709	C-29059	58	12/19/72 - 08/15/76 Antarctic Brightness

Monthly, Yearly, Multi-Year & 4-Yea



Technical Memorandum 83812

**Satellite-Derived Ice Data Sets
No. 1: Antarctic Monthly Average
Microwave Brightness
Temperatures and Sea Ice
Concentrations 1973-1976**

**H. J. Zwally, J. C. Comiso
and C. L. Parkinson**

NOVEMBER 1981

National Aeronautics and
Space Administration

Goddard Space Flight Center
Greenbelt, Maryland 20771

SATELLITE-DERIVED ICE DATA SETS NO. 1: ANTARCTIC
MONTHLY AVERAGE MICROWAVE BRIGHTNESS TEMPERATURES
AND SEA ICE CONCENTRATIONS, 1973-1976

H. J. Zwally, J. C. Comiso, and C. L. Parkinson
Goddard Laboratory for Atmospheric Sciences

November 1981

NASA/GODDARD SPACE FLIGHT CENTER
Greenbelt, Maryland 20771

SATELLITE-DERIVED ICE DATA SETS NO. 1: ANTARCTIC MONTHLY
AVERAGE MICROWAVE BRIGHTNESS TEMPERATURES AND SEA-ICE
CONCENTRATIONS, 1973-1976

H. J. Zwally, J. C. Comiso, and C. L. Parkinson
Goddard Laboratory for Atmospheric Sciences

ABSTRACT

A summary data set concerning 4 years of Antarctic sea ice conditions has been created and is available on magnetic tape¹. The data were derived from electrically scanning microwave radiometer (ESMR) brightness temperatures and were mapped into a polar stereographic grid enclosing the 50°S latitude circle. The grid size varies from about 32 km by 32 km at the poles to about 28 km by 28 km at 50°S. The microwave brightness temperatures of Antarctic sea ice are predominantly characteristic of first-year ice with an emissivity of 0.92 at 19 GHz frequency. Sea ice concentrations were calculated from the brightness temperature data for each grid element with an algorithm that uses an emissivity value of 0.92 and an ice physical temperature estimate from climatological surface air temperatures. Monthly, multiyear monthly, and yearly maps of brightness temperatures and sea ice concentrations were created for the 4 years, except for 7 months for which useable data were insufficient.

¹ The data are being released prior to the publication of a detailed analysis in an atlas entitled "Antarctic Sea Ice 1973-1976 from Satellite Passive Microwave Observations" by H. J. Zwally, J. C. Comiso, C. L. Parkinson, W. J. Campbell, F. D. Carsey, and P. Gloersen. Further analysis and application of the data are encouraged and any publication resulting from such study should reference the atlas as well as this document.

SATELLITE-DERIVED ICE DATA SETS NO. 1: ANTARCTIC
MONTHLY AVERAGE MICROWAVE BRIGHTNESS TEMPERATURES
AND SEA ICE CONCENTRATIONS, 1973-1976

H. J. Zwally, J. C. Comiso, and C. L. Parkinson
Goddard Laboratory for Atmospheric Sciences
NASA/Goddard Space Flight Center
Greenbelt, Maryland

INTRODUCTION

This report describes a summary data set concerning 4 years of Antarctic sea ice conditions, available on magnetic tape. The data set provides the most extensive available synoptic coverage of Antarctic sea ice, coverage made possible by the 19-GHz ESMR on board the Nimbus 5 satellite launched in December 1972. A detailed description of both the satellite and the instrument can be found in the Nimbus 5 User's Guide (Sabatini, 1972). The unique day/night capabilities of such radiometers and the strong contrast between the brightness temperature, T_B , of the ice and that of water at the 19-GHz frequency make the ESMR instrument suitable for observing spatial and temporal variations of sea-ice cover. The same data will appear in an Antarctic sea-ice atlas (Zwally et al., in preparation) in various forms, including false-color images of T_B at 5 K intervals, ice concentration at 4-percent intervals, and monthly differences in ice concentration at 5-percent intervals. In addition, the ice concentration maps will be presented at larger intervals as black-and-white images and as contour maps. The latter, which show ice concentration contours for 15-, 50-, and 85-percent ice concentration values, are included in this document as well.

The value of such sea-ice data sets for climate studies and for surface energy balance calculations has been cited by Fletcher (1968), Weller (1980), and Ackley (1981). In addition, Zwally and Gloersen (1977) have

noted other scientific applications. The commercial applicability of these data is also well recognized, particularly in areas in which oil exploration is actively pursued. Because of limited resolution (about 30 km by 30 km), however, the data are more useful for global or large-scale studies than for small-scale studies.

DATA PROCESSING, MAPPING, AND COMPILATION

The primary source of orbital Earth-located and calibrated radiometer data is the set of calibrated brightness temperature (CBT) tapes. The CBT tapes contain the time, calibration parameters, measured brightness temperatures, and corresponding geographical coordinates (Wilheit, 1972). The tapes are structured so that, when written at 6250 bits per inch, each tape contains about 180 orbits of data stored in separate files. Table 1 lists the format of each file in the CBT tape. These tapes were written in binary format and were blocked so that each block contains 50 records with 560 bytes per record.

To provide synoptic representation of the data in the polar regions for both spatial and temporal studies, the orbital data were projected to polar stereographic maps and accumulated at fixed time intervals. A polar stereographic map is a projection of points on the Earth's surface onto a plane tangent to the surface at a pole (north or south) with the vertex of the projection being the opposite pole. Figure 1 illustrates the concept, with a point (B) on the Earth's surface being projected to the polar plane at C through a line from the vertex pole (A) to B. The grid size for these maps is 293 by 293, with each map cell representing an area ranging from about 32 km by 32 km near the poles to about 28 km by 28 km near 50° latitude. The geographical coordinates of the center of the radiometer

field-of-view determine the cell into which the data is placed. Overlapping data in a cell from successive orbits are averaged to give a single brightness temperature average taken to be located at the center of the cell.

The coordinates of each point in the projected map are determined as follows. The distance (r) from the Pole to the projected point is calculated from:

$$r = d \tan \alpha \quad (1)$$

where d is the diameter of the Earth, and α is the angular projection at the opposite pole expressed in terms of the latitude ϕ as follows:

$$\alpha = \frac{90 - \phi}{2} . \quad (2)$$

With λ the east longitude and the Greenwich meridian coinciding with the positive ordinate of the map, the x and y components of the data element using Figure 1 are given by:

$$x = d \tan \alpha \cos (90^\circ - \lambda) \quad (3)$$

$$y = d \tan \alpha \sin (90^\circ - \lambda) \quad (4)$$

The standard 293 by 293-grid maps are constructed from the foregoing coordinate system by defining the location of each map cell by the following set of coordinates (I, J):

$$I = 147 + x + 0.5 \quad (5)$$

$$J = 147 - y + 0.5 \quad (6)$$

The value of 0.5 included in equations 5 and 6 properly positions the map cell at the center of its geographic location.

Full coverage of the entire polar area could be obtained from a sequence of six satellite orbits or one-half day of good data if the full set of 78 beam positions is used. However, because of the large disparity in

the radiometer field of view from the outer beam position to the middle beam position (i.e., 70 km by 140 km compared to 25 km by 25 km), only the middle 52 beam positions were used equatorward of 85 degrees latitude. This method provides a swath-angle coverage of $\pm 30.5^\circ$, corresponding to a spatial coverage of about 1280 km on the Earth's surface, and a minimum resolution of 29 km by 42 km. Because the area poleward of 85° latitude is observed by the outer beam positions only, the outer positions were used to cover this region. With this restriction on beam positions and with additional rejection of data because of instrumental problems, approximately 3 days of data are usually required for providing adequate spatial coverage of the polar regions. Three-day average maps were therefore generated for all periods, from launch until May 1977, that contained valid data.

Each data element on the polar stereographic maps represents a quantity (e.g., brightness temperature) averaged over the number of measurements made in that area within the specified time period. For a 3-day average map, this number of measurements, called population, ranges from 0 to 35, depending on the location of the data element and the availability of good data. During the analysis of the data, this value was used to determine how representative the data-point measurement is as an average over the given time. For most of the polar maps, a record of population normally follows a data record.

Because of data loss caused by processing and calibration problems, as many as 20 percent of the data cells were empty on some 3-day average maps. The 20 percent data loss occurred much more frequently after Nimbus 6 was launched in June 1975, when Nimbus 5 data became available only

every other day. Fortunately, most of these empty cells are located outside the range of the seasonal sea-ice extent. For completeness and proper weighting of data in the monthly average maps, some interpolations were made to fill in the empty cells in the 3-day average maps. A combination of temporal and spatial interpolation was employed. Normally, maps with very few empty cells were filled by linear spatial interpolation. For gaps involving four or more cells, however, time interpolation gave a more consistently satisfactory result. To fill in an empty cell by time interpolation, the average was taken of the brightness temperature in the cell for the previous 3-day average map and the one immediately after. In the rare cases when necessary, the procedure was extended to use data for up to three 3-day averages before and after the empty cell.

The brightness temperature of the ocean is expected to have minimal or no seasonal dependence, because the emissivity is inversely proportional to the surface physical temperature (Wilheit, 1972). Although brightness temperature is affected by roughness, foam cover, water vapor, and rainfall, none of these have seasonal characteristics. However, studies of the temporal variation of T_B in the southern ocean revealed some unexpected shifts; the conclusion was that these could be caused only by calibration or instrumental problems. Appropriate adjustments in T_B were made as follows.

To investigate the time dependence of T_B , the data for each pixel in the monthly maps between 55° and 65°S were binned and plotted. The contribution from the ocean appeared as a narrow peak with an average width of about ± 4 K. Figure 2 shows the brightness temperatures of these peaks for

each month of available ESMR data. In 1973, the values dropped by approximately 8 K from January to July, and, in 1976, a much sharper dip occurred in May, followed by a recovery in July and a general decrease for the remainder of the year. During 1974 and 1975, T_B was nearly constant with time. The apparently anomalous shift in T_B in 1976 was further studied by examining both ocean and highly concentrated sea-ice areas and using 3-day average maps. The shifts in brightness temperature in both ocean and ice areas occurred at about the same time. Furthermore, although the initial downward shift is about 10 K for the ocean and 8 K for sea ice, the recovery shift is about 18 K for the ocean and 25 K for sea ice. The latter shift would require a change in physical temperature of about 27 K on the ice and 40 K in the water, assuming emissivity to be constant during the time period. Because of the abrupt nature of the shifts and their magnitude, it is not possible to attribute them to physical effects in the ocean or the atmosphere.

Examination of the time variation of T_B of the Antarctic ice sheets also indicated that the smaller dip observed in 1973 resulted from calibration problems. The short-term year-to-year variation of T_B in the ice sheets was expected to be minimal because the seasonal physical temperature--the most significant variable expected to cause such a variation--did not change much during the 4-year period. By adjusting to force the ocean temperatures to remain constant, the T_B in the ice sheets also became more consistent during the 4 years. Comiso and Zwally (1980) discuss the procedures used for normalizing the data in response to these anomalous shifts.

EXTRACTION OF SEA-ICE CONCENTRATION FROM ESMR

Within the ice pack, the brightness temperature of a pixel area derives from various sources--the water and ice within the field of view, and the atmosphere. To a good approximation,

$$T_B = C_W \epsilon_W T_W + C_I \epsilon_I T_I + T_A \quad (7)$$

where ϵ_W , T_W , and C_W are emissivity, surface physical temperature, and areal percentage of water, and ϵ_I , T_I , and C_I are the corresponding values for sea ice. T_A is the contribution attributable to the atmosphere and includes the direct upwelling radiation, the downwelling radiation reflected from the surface, and the radiation from space reflected by the surface. Because the reflectance of water is different from that of ice, the atmospheric contribution can be separated into contributions over water, T_{AW} , and over ice, T_{AI} (i.e., $T_A = C_W T_{AW} + C_I T_{AI}$). The atmospheric opacity in the polar region has been estimated to be very small and was neglected. Also, $C_W + C_I = 1$ and ϵ_I was determined empirically to be about 0.92.

The ice concentration, C_I , can thus be determined from equation 7:

$$C_I = \frac{T_B - T_0}{\epsilon_I T_{\text{eff}} - T_0} \quad (8)$$

where $T_0 = \epsilon_W T_W + T_{AW}$ is the measured brightness temperature of the water determined from the data, and $T_{\text{eff}} = T_I + T_{AI}/\epsilon_I$ is the effective surface physical temperature. The value of T_{AW} was calculated by using radiative transfer modeling of the atmosphere (Nieman and Wilheit, private communication) and was estimated over the ocean to be about 8 K. The value of T_{AI} is considerably smaller because the reflectivity of ice is substantially lower than that of water, and the contribution of downwelling

radiation reflected from the surface is therefore smaller. In the polar regions, the atmospheric contribution to the brightness temperature is minimal because the humidity and the water-vapor content in the atmosphere is very low. Therefore, T_{eff} is taken to be equal to T_I . The appropriate physical temperature, T_I , is the temperature at the top of the sea ice below the snow cover, because most of the observed radiation emanates from a thin top layer of saline ice. In the absence of real-time physical temperature data, T_I was estimated from a compilation of climatological surface air temperatures (Jenne et al., 1974) as follows. The actual temperature of the top sea-ice surface lies between the surface air temperature, T_{air} , interpolated from the climatological data, and the temperature of the water underneath the ice. It can be expressed by:

$$T_I = T_{\text{air}} + f(T_m - T_{\text{air}}) \quad (9)$$

where T_m is the melting temperature of ice, and f is an empirical parameter deduced from the ESMR data. Figure 3 shows a histogram of the ice-covered area in the Weddell Sea in July 1973 as a function of ice concentration. The parameter f was adjusted until the backward peak of the histogram fell off approximately near the 100-percent region, allowing for several degrees of uncertainty in the observed brightness temperatures. The value of $f = 0.25$ obtained by this procedure gives temperatures of the snow/ice interface relative to T_{air} , which are in good agreement with surface measurements on first-year ice in the Arctic (Ramseier, private communication). The value of f will vary spatially with the thicknesses of the ice and the snow cover, but $f = 0.25$ appears to be a good average value. Also, analysis of the open-ocean area in the vicinity of the ice pack showed that $T_0 \cong 135$ K. Thus, two reference points were used in determining ice concentration.

DATA STORAGE AND RETRIEVAL

The 4-year data sets of southern ocean ESMR brightness temperatures, plus the sea ice concentrations calculated with the foregoing equations and considerations, are stored in sequential arrays on magnetic tape. Each polar-map cell is associated with a matrix element, $D(J,I)$, that represents the average value of the data over the number of measurements represented by the population matrix, $P(J,I)$, where J and I are as defined in equations 5 and 6. The sequence in which the data are stored is shown in Table 2. A 20-word, 4 byte-per-word header preceding each map describes the type of map, the size of the grid, the scaling, the time interval the map data were accumulated, and other information associated with the data (Table 3). This header is followed by a sequence of 293 times 2 data and population records, each 293 words long representing $D(J,I)$ and $P(J,I)$ for constant I and for J varying from 1 to 293. For the data and population records, each word is 5 bytes, making each record 1465 bytes. Note that on the magnetic tapes, in order to maintain data precision along with an integer format, each data word has been multiplied by a scaling factor of 50. Although only the first 80 bytes are meaningful on the header record, this record also has a length of 1465 bytes. The data are blocked when written on tape so that there are 5 records per block. A sample Fortran program and job control language for reading the tape in an IBM computer facility is given in the Appendix.

All the monthly data have been placed on one magnetic tape written at 6250 bits per inch. The 41 matrices of monthly T_B 's from 1973 to 1976 are stored in files 1 through 41 in chronological order, followed by 12 matrices of multiyear monthly T_B 's and 5 matrices of yearly and 4-year average T_B 's. These are followed by the corresponding 58 ice-concentration matrices, the 12 climatological surface air temperature matrices, and the 12 climatological pressure matrices. Table 4 summarizes the population and overall quality

of each polar map. Figures 4 to 13 show contour maps of monthly and multiyear monthly ice concentrations.

Three values of ice concentration were contoured: 15, 50, and 85 percent. A detailed analysis of the data set and descriptions of the ice conditions are given in the aforementioned atlas (Zwally et al., in preparation) which includes pseudo color images of brightness temperature, sea ice concentration, changes in ice concentration, and multiyear average concentrations. Also included are graphs of the area of ice in various concentration categories as a function of time for the entire southern ocean and in five geographical sectors. The amount of open water, mean ice concentration, and actual ice areas are shown. The atlas further includes discussions of the observed interannual variations and the characteristics of the growth-decay cycle.

ACKNOWLEDGEMENTS

The authors wish to thank M. Osborne, R. Pelletier, and D. Love from Computer Sciences Corporation, and J. Manning and D. Johnson from Computer Sciences Technicolor Associates for programming and technical support in the preparation of the data sets. We also appreciate discussions on these data sets with Drs. P. Gloersen, W. J. Campbell, F. D. Carsey and R. Ramseier. This work was supported by NASA's Climate Program.

REFERENCES

- Ackley S. F., "A Review of Sea-ice Weather Relationships in the Southern Hemisphere," Sea Level, Ice and Climatic Change (Proceedings of the Canberra Symposium, December 1979). IAHS Publication 131, 1981.
- Comiso, J. C., and H. J. Zwally, "Corrections for Anomalous Time Dependent Shifts in the Brightness Temperature from the Nimbus 5 ESMR," NASA TM-82055, 1980.
- Fletcher, J. O., 1968: The influence of the Arctic pack ice on climate. In: Meteorological Monographs, J. M. Mitchell, Jr., ed., (Am. Met. Society), 8, 93-99.
- Jenne, R. L., H. L. Crutcher, H. Van Loon, and J. J. Taljaard, "A Selected Climatology of the Southern Hemisphere: Computer Methods and Data Availability," NCAR-TN/STR-92, 1974.
- Sabatini, R. R., ed., Nimbus 5 User's Guide, NASA/Goddard Space Flight Center, 1972.
- Weller, G., "Spatial and Temporal Variations in the South Polar Surface Energy Balance," Monthly Weather Review, 108, 1980, pp.2006-2014.
- Wilheit, T. T., "The Electrically Scanning Microwave Radiometer (ESMR) Experiment," Nimbus 5 User's Guide, NASA/Goddard Space Flight Center, 1972, pp. 59-105.
- Zwally, H. J., J. C. Comiso, C. L. Parkinson, W. J. Campbell, F. D. Carsey, and P. Gloersen, "Antarctic Sea Ice Cover 1973-1976 from Satellite Passive Microwave Observations," NASA SP, in preparation.
- Zwally, H. J., and P. Gloersen, "Passive Microwave Images of the Polar Regions and Research Applications," Polar Rec., 18, 1977, pp.431-450.

Table 1
CBT Data Record Format for ESMR

Word	Quantity	Units	Scale	Description
1	Year	Year		Year associated with data
2	Day	Days		
3	Hour	Hours		
4	Minute	Minutes		
5	Second	Seconds		
6	Program ID			Unique Program Identification
7	Pitch error	Degrees	x10	Pitch fine error
8	Roll error	Degrees	x10	Roll fine error
9	RMP		x10	
	Indicated			
	Rate high			
10	Latitude	Degrees	x10	Latitude of subsatellite point
11	Longitude	Degrees	x10	Longitude of subsatellite point
12	Height	Kilometers		Height of spacecraft
13	Hot-load mean		x10	
14	Hot load		x100	
15	Cold-load mean		x10	
16	Cold load		x100	
17	MUX 1			Average antenna temperature
18	MUX 2			Phase-shift temperature
19	MUX 3			Ferrite-switch temperature
20	MUX 4			Ambient load temperature
21	MUX 5			Hot-load temperature
22	MUX 6			Automatic gain control
23-41	Engineering data			
42	Beam			
	Position 79			
43-46	MUX 1-MUX 4			
47-124	Latitude	Degrees	x10	Latitudes of the 78 scan positions
125-202	Longitude	Degrees	x10	Longitudes of the 78 scan positions
203-280	Brightness temperature	Kelvins	x10	Brightness temperatures of the 78 scan positions

Source: Nimbus 5 User's Guide, 1972.

Table 2

Standard Format* for Polar Stereographic Maps

Record Number	Length (bytes)	Format	Number of Words	Description
1	1465	(see below) ⁺	293	Heading (only the first 20 words are meaningful)
2	1465	293I5	293	Row 1 data (x50)
3	1465	293I5	293	Observation population of row 1
4	1465	293I5	293	Row 2 data (x50)
5	1465	293I5	293	Observation population of row 2
.	.	.	.	.
.	.	.	.	.
.	.	.	.	.
586	1465	293I5	293	Row 293 data (x50)
587	1465	293I5	293	Observation population of row 293

*Data format 2 indicates standard format (see Table 3).
⁺(9I5, 2A5, 8I5, A5, 255I5, 18I5)

Table 3
Header Record for Polar Stereographic Maps

Word	Type	Number of bytes/words	Description
1	I5	4	Projection type (1 for polar stereographic)
2	I5	4	Number of columns (293)
3	I5	4	Number of rows (293)
4	I5	4	Scale (2.5 x 106)
5	I5	4	Latitude enclosed (-50.0°)
6	I5	4	Greenwich orientation (270°)
7	I5	4	Radius of the Earth
8	I5	4	J-coordinate of the pole (147)
9	I5	4	I-coordinate of the pole (147)
10-11	A5	4	Data type* (TB, ICE CON, SURF TEMP, or CLIM PRES)
12	I5	4	Start time (day)
13	I5	4	Start time (hour)
14	I5	4	Start time (minutes)
15	I5	4	Stop time (day)
16	I5	4	Stop time (hour)
17	I5	4	Stop time (minutes)
18	I5	4	Year data was collected
19	I5	4	Data format (2 for standard format)
20	A5	4	Data Identification
21-293	I5	4	Meaningless

* Variable abbreviations for words 10 and 11 are as follows: TB is brightness temperature, ICE CON is ice concentration, SURF TEMP is mean monthly climatological surface air temperature, and CLIM PRES is mean monthly climatological sea level pressure.

Table 4

Nimbus 5 Ice Data Set

Date	Time Interval (Julian Date)	Population	File No.	Data Quality
<u>1. ESMR Brightness Temperature Maps</u>				
January 1973	001-031	238	1	Good
February 1973	032-059	287	2	Good
June 1973	152-181	342	3	Good
July 1973	182-212	271	4	Good
September 1973	249-273	127	5	Fair
October 1973	274-304	270	6	Good
November 1973	305-332	261	7	Good
December 1973	336-265	211	8	Good
January 1974	001-031	263	9	Good
February 1974	032-059	305	10	Good
March 1974	060-081	247	11	Good
April 1974	097-120	260	12	Good
May 1974	121-151	274	13	Good
June 1974	152-181	296	14	Good
July 1974	182-212	222	15	Good
August 1974	213-243	203	16	Good
September 1974	244-273	321	17	Good
October 1974	274-304	256	18	Good
November 1974	305-334	275	19	Good
December 1974	335-364	323	20	Good
January 1975	001-028	333	21	Good
February 1975	032-059	269	22	Good
March 1975	060-088	298	23	Good
April 1975	098-106	69	24	Poor
May 1975	122-151	325	25	Good
September 1975	244-273	62	26	Poor
October 1975	274-304	124	27	Fair
November 1975	305-334	112	28	Fair
December 1975	335-365	157	29	Fair
January 1976	001-031	116	30	Fair
February 1976	032-060	154	31	Fair
March 1976	061-091	146	32	Fair
April 1976	092-121	179	33	Fair
May 1976	122-152	151	34	Fair
June 1976	153-182	148	35	Fair
July 1976	186-213	155	36	Fair
August 1976	214-244	189	37	Fair
September 1976	245-274	163	38	Fair
October 1976	275-305	173	39	Fair
November 1976	306-335	169	40	Fair
December 1976	336-365	104	41	Fair

Table 4
Nimbus 5 Ice Data Sets (continued)

Date	Time Interval (Julian Date)	Total Orbits	File No.	Data Quality
January 1973-76	001-031	993	42	Good
February 1973-76	032-060	1015	43	Good
March 1974-76	061-091	691	44	Good
April 1974-76	092-121	508	45	Fair
May 1974-76	122-152	750	46	Good
June 1973/74/76	153-182	786	47	Good
July 1973/74/76	183-213	648	48	Good
August 1974/76	214-244	392	49	Fair
September 1973-76	245-274	673	50	Good
October 1973-76	275-305	823	51	Good
November 1973-76	306-335	817	52	Good
December 1973-76	336-365	795	53	Good
1973	001-365	2050	54	Fair
1974	001-365	3245	55	Good
1975	001-365	1749	56	Fair
1976	001-366	1847	57	Good
1973-76	001-365	8891	58	Good

2. Sea Ice Concentration Monthly Maps

January 1973	001-031	281	59	Good
February 1973	032-059	287	60	Good
June 1973	152-181	342	61	Good
July 1973	182-212	271	62	Good
September 1973	249-273	127	63	Fair
October 1973	274-304	270	64	Good
November 1973	305-332	261	65	Good
December 1973	336-265	211	66	Good
January 1974	001--31	263	67	Good
February 1974	032-059	305	68	Good
March 1974	060-081	247	69	Good
April 1974	097-120	260	70	Good
May 1974	121-151	274	71	Good
June 1974	152-181	296	72	Good
July 1974	182-212	222	73	Good
August 1974	213-243	203	74	Good
September 1974	244-273	321	75	Good
October 1974	274-304	256	76	Good
November 1974	305-334	275	77	Good
December 1974	335-364	322	78	Good
January 1975	001-028	333	79	Good
February 1975	032-059	269	80	Good
March 1975	060-088	298	81	Good
April 1975	098-106	69	82	Poor
May 1975	122-151	325	83	Good
September 1975	244-273	62	84	Poor

Table 4
Nimbus 5 Ice Data Sets (continued)

Date	Time Interval (Julian Date)	Total Orbits	File No.	Data Quality
October 1975	274-304	124	85	Fair
November 1975	305-334	112	86	Fair
December 1975	335-365	157	87	Fair
January 1976	001-031	116	88	Fair
February 1976	032-060	154	89	Fair
March 1976	061-091	146	90	Fair
April 1976	092-121	179	91	Fair
May 1976	122-152	151	92	Fair
June 1976	153-182	148	93	Fair
July 1976	186-213	155	94	Fair
August 1976	214-244	189	95	Fair
September 1976	245-274	163	96	Fair
October 1976	275-305	173	97	Fair
November 1976	306-335	169	98	Fair
December 1976	336-365	104	99	Fair
January 1973-6	001-031	993	100	Good
February 1973-6	032-060	1015	101	Good
March 1974-6	061-091	691	102	Good
April 1974-6	092-121	508	103	Fair
May 1974-6	122-152	750	104	Good
June 1973,74,76	153-182	786	105	Good
July 1973,74,76	183-213	648	106	Good
August 1974,76	214-244	392	107	Fair
September 1973-6	245-274	673	108	Good
October 1973-6	275-305	823	109	Good
November 1973-6	306-335	817	110	Good
December 1973-6	336-365	795	111	Good
1973	001-365	2050	112	Fair
1974	001-365	3245	113	Good
1975	001-365	1749	114	Fair
1976	001-366	1847	115	Good
1973-6	001-365	8891	116	Good

3. Monthly Climatological Surface Temperature Maps

January	001-031	117
February	032-059	118
March	060-081	119
April	097-120	120
May	121-151	121
June	152-181	122
July	182-212	123
August	213-243	124
September	244-273	125

Table 4
Nimbus 5 Ice Data Sets (continued)

Date	Time Interval (Julian Date)	Total Orbits	File No.	Data Quality
October	274-304		126	
November	305-334		127	
December	335-364		128	

4. Monthly Climatological Sea-Level Pressure Maps

January	001-031	129
February	032-059	130
March	060-081	131
April	097-120	132
May	121-151	133
June	152-181	134
July	182-212	135
August	213-243	136
September	244-273	137
October	274-304	138
November	305-334	139
December	335-364	140





Figure 1. Schematic Diagram for Polar Stereographic Mapping.




Figure 2. Mean brightness temperatures of the ocean from monthly average maps for 1973-1976 before correction for calibration shifts.




Figure 3. Ice concentration distribution near the Weddell Sea in July 1973.

ICE CONCENTRATION BOUNDARIES FOR 1973




Figure 4. Monthly ice concentration contours for January through June 1973.

ICE CONCENTRATION BOUNDARIES FOR 1973




Figure 5. Monthly ice concentration contours for July through December 1973.

ICE CONCENTRATION BOUNDARIES FOR 1974




Figure 6. Monthly ice concentration contours for January through June 1974.

ICE CONCENTRATION BOUNDARIES FOR 1974

—— 15.0% BOUNDARY 50.0% BOUNDARY ===== 85.0% BOUNDARY




Figure 7. Monthly ice concentration contours for July through December 1974.

ICE CONCENTRATION BOUNDARIES FOR 1975




Figure 8. Monthly ice concentration contours for January through June 1975.

ICE CONCENTRATION BOUNDARIES FOR 1975

—— 15.0% BOUNDARY 50.0% BOUNDARY ||||| 85.0% BOUNDARY




Figure 9. Monthly ice concentration contours for July through December 1975.

ICE CONCENTRATION BOUNDARIES FOR 1976




Figure 10. Monthly ice concentration contours for January through June 1976.

ICE CONCENTRATION BOUNDARIES FOR 1976




Figure 11. Monthly ice concentration contours for July through December 1976.

ICE CONCENTRATION BOUNDARIES

—— 15.0% BOUNDARY 50.0% BOUNDARY ===== 85.0% BOUNDARY




Figure 12. Multiyear monthly ice concentrations contours for January through June 1973-1976.

ICE CONCENTRATION BOUNDARIES

—— 15.0% BOUNDARY 50.0% BOUNDARY ===== 85.0% BOUNDARY




Figure 13. Multiyear monthly ice concentrations contours for July through December 1973-1976.

APPENDIX

I. Sample Fortran program for reading the data tape:

```
DIMENSION IHEAD (293), IDAT(293), IPOP(293)
```

```
.  
.  
.
```

```
READ (8,510) IHEAD
```

```
.  
.  
.
```

C

```
DO 50 J=1, 293
```

```
READ (8,520) IDAT
```

```
READ (8, 520) IPOP
```

C

```
50 CONTINUE
```

```
510 FORMAT (9I5, 2A5, 8I5, A5, 255I5, 8I5)
```

```
520 FORMAT (255I5, 38I5)
```

```
END
```

II. Job control language tape density and blocking specification (IBM System):

```
//FT08F001 DD UNIT = (6250, DEFER), LABEL = (1,NL), DISP = OLD,
```

```
// DCB = (DEN = 4, RECFM = FB, LRECL = 1465, BLKSIZE = 7325)
```

BIBLIOGRAPHIC DATA SHEET

1. Report No. TM-83812	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Satellite-Derived Ice Data Sets No.1: Antarctic Monthly Average Microwave Brightness Temperatures and Sea-Ice Concentrations, 1973-1976.		5. Report Date November 1981	
		6. Performing Organization Code 912.1	
7. Author(s) H. J. Zwally, J. C. Comiso, C. L. Parkinson		8. Performing Organization Report No.	
9. Performing Organization Name and Address Goddard Space Flight Center Greenbelt, MD 20771		10. Work Unit No.	
		11. Contract or Grant No.	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, DC 20546		13. Type of Report and Period Covered Technical Memorandum	
		14. Sponsoring Agency Code	
15. Supplementary Notes			
16. Abstract A summary data set concerning 4 years of Antarctic sea ice conditions has been created and is available on magnetic tape¹. The data were derived from electrically scanning microwave radiometer (ESMR) brightness temperatures and were mapped into a polar stereographic grid enclosing the 50°S latitude circle. The grid size varies from about 32 km by 32 km at the poles to about 28 km by 28 km at 50°S. The microwave brightness temperatures of Antarctic sea ice are predominantly characteristic of first-year ice with an emissivity of 0.92 at 19 GHz frequency. Sea ice concentrations were calculated from the brightness temperature data for each grid element with an algorithm that uses an emissivity value of 0.92 and an ice physical temperature estimate from climatological surface air temperatures. Monthly, multiyear monthly, and yearly maps of brightness temperatures and sea ice concentrations were created for the 4 years, except for 7 months for which useable data were insufficient. ¹ The data are being released prior to the publication of a detailed analysis in an atlas entitled "Antarctic Sea Ice 1973-1976 from Satellite Passive Microwave Observations" by H. J. Zwally, J. C. Comiso, C. L. Parkinson, W. J. Campbell, F. D. Carsey, and P. Gloersen. Further analysis and application of the data are encouraged and any publication resulting from such study should reference the atlas as well as this document.			
17. Key Words (Selected by Author(s)) Sea Ice Data, Microwave Brightness, Antarctic		18. Distribution Statement STAR Category Unclassified-Unlimited	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 34	22. Price* \$4.50

D 47329

11/173 - 12/13/176

201

NTN

ATLS

```
RECCFD      = 1 OF FILE
LENGTH     = 7325 BYTES
```

[illegible]

40F7F2F6 F640F7F2 F6F040F7 F4F3F640 F7F3F1F4 40F7F3F4 F340F7F3 F6F440F7 F4F6F140 F7F3F4F0
40F7F5F2 F340F7F4 F0F64040 4CF5F640 4040F6F7 404040F6 F34040F1 F0F04040 40F6F540 4040F9F7
404040F6 F5404040 F5F64040 4CF6F240 40F1F5F2 4040F1F0 F1404040 F9F54040 40F7F240 4040F6F2
404CF1F0 F6404040 F6F74040 F1F1F540 4040F6F1 404040F7 F04040F1 F4F44040 40F6F540 4040F5F9
404040F7 F04040F1 F5F74040 4CF6F640 40F1F4F0 404040F6 F2404040 F6F54040 40F6F740 4040F6F0
404040F6 F6404040 F2F54040 4CF6F540 40F1F8F4 404040F6 F4404040 F2F44040 40F2F540 4040F2F8
4040F1F3 F6404040 F2F74040 4CF6F740 4040F2F2 404040F6 F5404040 F5F64040 40F6F640 4040F2F8
404040F3 F0404040 F6F24040 4CF2F640 4040F2F3 404040F6 F2404040 F2F74040 40F5F540 4040F6F4
404040F6 F1404040 F6F04040 F1F0F340 40F1F1F0 404040F6 F2404040 F6F04040 F1F0F140 4040F2F2
404040F2 F6404040 F2F54040 F1F6F240 4040F2F4 404040F6 F1404040 F2F64040 40F2F540 4040F2F9
404040F3 F0404040 F2F44040 4CF2F740 4040F2F8 404040F2 F1404040 F2F54040 4CF3F140 4040F6F5
404040F2 F5404040 F6F54040 4CF2F140 4040F2F8 404040F1 F5404040 F2F64040 F1F1F440 4040F3F0
404040F5 F5404040 F2F64040 4CF6F340 4040F6F8 404040F6 F5404040 F5F74040 F1F0F340 4040F6F3
404CF1F0 F1404040 F2F54040 4CF7F340 4040F2F2 404040F3 F0404040 F6F64040 4CF3F140 4040F6F1
404040F6 F1404040 F2F54040 4CF2F140 4040F6F8 404040F2 F64040F1 F0F54040 40F6F440 4040F7F3
404040F6 F1404040 F7F44040 4CF7F540 4040F7F2 404040F1 F6404040 F6F24040 4CF8F140 4040F3F2
404040F4 F0404040 F7F04040 4CF6F640 4040F3F8 404040F3 F3404040 40F2F640 4040F3F3
404040F3 F7404040 F6F44040 4CF3F540 4040F4F5 404040F2 F5404040 F3F34040 40F4F240 4040F8F3
404040F2 F74040F1 F4F04040 4CF3F340 4040F5F1 404040F4 F5404040 F4F34040 40F3F040 4040F2F8
404040F3 F2404040 F3F34040 4CF6F540 4040F3F0 404040F2 F4404040 F6F74040 40F3F740 4040F3F4
404040F3 F0404040 F3F14040 4CF3F540 4040F4F8 404040F5 F1404040 F5F24040 40F3F640 4040F2F7
404040F3 F3404040 F4F64040 4CF4F740 4040F4F0 404040F4 F2404040 F5F04040 40F4F440 4040F3F6
404040F3 F6404040 F4F24040 4CF5F540 4040F4F8 404040F7 F5404040 F3F54040 40F4F440 4040F4F0
404040F4 F5404040 F7F64040 4CF5F640 4040F4F7 404040F8 F1404040 F8F34040 40F5F140 4040F8F2
404040F7 F7404040 F8F84040 4CF8F240 4040F8F6 404040F5 F1404040 F8F14040 40F8F140 4040F5F3
404040F7 F5404040 F3F64040 4CF4F740 4040F5F2 404040F4 F2404040 F3F54040 40F5F240 4040F4F5
404040F4 F3404040 4CF4F240 4040F3F5 404040F3 F6404040 F3F54040 40F4F540 4040F5F2
404040F3 F4404040 F4F34040 4CF3F640 4040F4F5 404040F3 F7404040 F4F34040 40F4F040 4040F4F9
404040F3 F6404040 F4F54040 4CF4F340 4040F4F0 404040F4 F4404040 F5F54040 40F3F340 4040F4F1
404040F4 F0404040 F5F04040 4CF4F040 4040F4F3 404040F4 F3404040 F3F84040 40F3F940 4040F4F0
404040F4 F6404040 F4F14040 4CF3F740 4040F3F6 404040F3 F7404040 F3F54040 40F3F640 4040F4F2
404040F4 F1404040 F3F64040 40F4F140 4040F3F7 404040F3 F3404040 F4F44040 40F3F240 4040F4F4
404040F4 F6404040 F4F24040 4CF3F740 4040F3F1 404040F4 F1404040 F4F14040 40F3F640 4040F3F5
404040F2 F5404040 F3F64040 4CF6F340 4040F2F6 404040F3 F74040F1 F0F34040 40F3F540 4040F3F2
404040F3 F6404040 F2F74040 40F3F740 4040F3F1 4040F1F0 F6404040 F3F34040 40F7F240 4040F3F4
404040F3 F5404040 F6F54040 4CF7F740 4040F3F8 404040F2 F5404040 F2F24040 40F3F040 4040F3F5
404040F6 F6404040 F6F54040 4CF7F740 4040F6F7 404040F7 F2404040 F6F54040 40F3F140 4040F5F7
404040F3 F74040F1 F0F14040 40F3F240 4040F2F3 404040F6 F1404040 F2F64040 40F2F240 F7F1F8F7
40F7F6F0 F440F7F7 F7F640F7 F4F6F440 F7F7F9F2 40F7F4F4 F640F7F2 F3F340F7 F0F7F440 F7F2F5F7
40F7F4F6 F640F7F1 F7F040F7 F2F1F640 F7F1F2F6 40F7F1F5 F440F7F2 F5F440F7 F1F2F540 F7F2F0F9
40F7F0F9 F340F7F3 F0F440F7 F1F5F740 F7F1F8F3 40F7F1F2 F440F7F0 F4F240F7 F0F6F640 F7F0F7F2
40F7F0F2 F440F7F0 F7F240F7 F6F0F140 F6F9F8F0 40F7F0F7 F340F7F0 F5F640F6 F5F9F340 F7F0F0F8
40F7F0F1 F740F6F9 F5F440F7 F0F4F540 F7F0F5F5 40F7F0F6 F640F6F5 F8F640F7 F0F2F540 F7F0F6F8
40F6F9F5 F540F6F9 F6F140F6 F0F4F140 F6F9F6F3 40F6F9F5 F440F6F5 F8F540F7 F0F4F140 F6F9F8F8
40F7F0F5 F040F6F9 F7F30F7 F0F5F540 F6F9F1F8 40F7F1F3 F540F6F9 F0F7F240 F0F7F240 F7F1F3F1
40F7F1F6 F140F7F0 F6F240F7 F0F6F240 F7F2F3F4 40F7F1F5 F640F7F0 F2F640F7 F1F6F640 F7F1F5F5
40F7F1F5 F340F7F3 F1F740F7 F1F6F640 F7F3F4F5 40F7F2F0 F240F7F2 F5F640F7 F3F3F040 F7F2F2F1
40F7F3F0 F540F7F2 F7F540F7 F1F6F640 F7F0F2F6 40F6F9F7 F1F640F7 F1F0F440 F6F5F8F9
40F6F5F5 F340F7F0 F6F540F6 F5F3F640 F7F1F2F3 40F6F5F4 F540F7F0 F4F940F6 F5F5F340 F6F9F1F5
40F6F6F4 F440F6F9 F1F740F6 F6F6F640 F6F9F7F5 40F7F1F0 F140F6F9 F0F140F6 F5F0F640 F6F9F5F3
40F7F0F5 F140F6F4 F2F340F7 F0F5F540 F7F1F8F4 40F7F1F5 F540F7F1 F4F740F7 F1F9F340 F6F5F5F4
40F7F0F2 F540F6F9 F3F140F6 F5F7F140 F6F9F3F1 40F6F6F5 F640F6F5 F8F640F7 F0F2F540 F6F9F5F3
40F7F0F4 F540F6F9 F7F040F6 F5F7F540 F6F9F8F2 40F6F5F2 F740F6F5 F5F540F6 F5F0F640 F6F8F8F7
40F6F5F4 F540F6F8 F5F740F6 F5F3F140 F6F5F4F5 40F7F0F1 F340F7F0 F6F640F6 F5F6F240 F6F9F6F1
40F7F0F2 F340F7F0 F2F840F6 F5F6F740 F7F0F2F1 40F7F0F5 F240F7F0 F3F340F6 F5F4F740 F6F8F8F4
40F6F5F3 F140F6F9 F1F240F6 F6F2F640 F6F8F6F3 40F6F5F1 F440F6F9 F1F140F6 F6F8F240 F6F5F7F1
40F6F6F6 F040F6F8 F6F240F6 F6F6F540 F6F9F2F8 40F6F5F1 F640F6F8 F6F240F6 F6F3F840 F6F5F5F5
40F6F6F5 F640F6F8 F7F640F6 F6F6F540 F6F8F5F5 40F6F6F5 F440F6F8 F9F540F6 F5F5F040 F6F5F4F9
40F6F9F0 F640F6F9 F6F340F7 F0F1F640 F6F9F5F2 40F6F5F3 F740F6F9 F0F340F6 F5F1F740 F6F8F5F1
40F6F9F3 F740F6F8 F6F340F7 F0F1F540 F6F8F4F5 40F6F9F1 F440F6F8 F1F040F6 F6F0F340 F6F8F7F3
40F6F8F0 F640F6F7 F4F540F6 F7F5F640 F6F7F5F1 40F6F6F7 F440F6F7 F7F040F6 F7F7F540 F6F8F3F1
40F6F8F3 F440F6F7 F5F740F6 F7F6F140 F6F8F3F5 40F6F6F7 F240F6F7 F8F240F6 F7F9F640 F6F8F3F6
40F6F8F5 F040F6F7 F6F540F6 F6F5F040 F6F8F3F5 40F6F8F0 F540F6F7 F7F640F6 F5F1F240 F6F8F8F7
40F6F8F4 F340F6F8 F4F640F6 F7F5F540 F6F9F0F0 40F6F8F6 F440F6F8 F5F640F6 F7F5F240 F6F8F6F3
40F6F8F1 F140F6F8 F2F740F6 F6F1F640 F6F8F8F9 40F6F8F6 F040F6F8 F8F540F6 F6F5F340 F6F9F2F3
40F6F8F3 F540F6F9 F0F640F6 F7F5F540 F6F7F8F8 40F6F8F0 F0F740F6 F6F2F140 F6F7F7F8
40F6F8F4 F140F6F8 F4F440F6 F6F1F240 F6F7F6F2 40F6F9F5 F1F740F6 F6F0F140 F6F6F5F8
40F6F7F4 F640F6F8 F1F040F6 F5F6F140 F6F9F1F6 40F6F8F6 F640F6F9 F1F640F6 F5F4F140 F6F8F6F3
40F6F5F7 F640F6F8 F8F540F7 F0F6F140 F6F8F6F5 40F6F5F0 F1F240F7 F0F8F540 F7F0F2F6
40F6F5F6 F140F6F9 F6F640F7 F1F1F540 F7F0F5F7 40F7F1F5 F740F7F2 F1F6F540 F7F0F0F5
40F7F1F6 F640F6F9 F5F440F7 F0F2F740 F7F0F6F4 40F7F0F0 F340F7F2 F0F540F7 F1F3F740 F7F1F7F6
40F7F0F2 F640F6F9 F1F540F7 F0F3F140 F7F0F5F2 40F7F0F3 F340F7F0 F2F640F7 F0F6F240 F7F1F8F0
40F7F0F6 F640F7F1 F1F340F7 F1F2F740 F6F9F6F7 40F7F1F6 F440F7F1 F2F7F540 F7F2F6F0
40F7F2F8 F140F7F2 F4F040F7 F1F5F540 F7F2F3F6 40F7F3F0 F640F7F3 F0F640F7 F4F2F040 F7F3F5F3
40F7F4F4 F640F7F3 F6F740F7 F5F1F140 F7F4F4F6 404040F6 F54040F1 F0F24040 4CF5F340 4040F7F4
404CF1F8 F5404040 F6F64040 4CF6F340 4040F1F4F5 404040F6 F34040F1 F4F24040 4CF6F440 40F1F1F3
404CF1F4 F2404040 F6F14040 4CF6F340 4040F6F6 4040F1F4 F6F44040 F6F44040 F1F4F040 4040F6F7
404040F6 F1404040 F6F14040 4CF6F540 4040F6F7 404040F2 F0404040 F9F84040 F1F0F040 40F1F0F5
404CF1F0 F0404040 F6F44040 4CF6F440 4040F6F3 404040F2 F5404040 F9F84040 40F2F740 4040F6F5
404040F5 F14040F1 F1F54040 4CF2F540 4040F6F5 404040F6 F2F54040 40F2F540 4040F2F6
404040F2 F34040F1 F0F24040 4CF1F640 4040F6F1 404040F2 F6404040 F2F04040 40F2F440 4040F6F1
404040F6 F5404040 F6F34040 F1F6F240 40F1F5F2 404040F6 F3404040 F6F24040 4CF1F840 40F1F0F5
404040F6 F1404040 F2F54040 4CF6F140 4040F6F6 404040F2 F6404040 F2F54040 40F2F540 4040F2F7

404040F5	F7404040	F3F44040	40F3F340	4040F2F2	404040F2	F5404040	F2F44040	40F3F240	4040F6F5
404040F6	F5404040	F2F64040	40F6F340	4040F6F6	404040F2	F5404040	F6F14040	40F2F640	4040F6F2
404040F2	F3404040	F6F14040	40F6F440	4040F6F0	404040F6	F74040F1	F5F44040	40F6F340	4040F6F8
404040F7	F4404040	F5F84040	40F2F640	4040F3F0	404040F3	F2404040	F2F94040	F1F1F740	4040F2F7
404040F2	F5404040	F6F84040	40F6F540	40F1F0F5	404040F3	F4404040	F7F24040	40F6F840	4040F3F3
404040F7	F7404040	F6F44040	40F7F540	4040F6F6	404040F7	F44040F1	F1F04040	40F7F040	4040F3F8
404040F6	F5404040	F3F74040	40F3F140	4040F4F3	404040F3	F3404040	F2F54040	40F4F840	4040F2F3
404040F3	F6404040	F4F44040	40F3F040	4040F4F5	404040F3	F2404040	F2F74040	40F3F640	4040F7F8
404040F3	F0404040	F2F94040	40F4F140	4040F4F6	404040F3	F5404040	F2F54040	40F2F840	4040F3F5
404040F4	F4404040	F4F54040	40F5F140	4040F5F1	404040F5	F0404040	F4F44040	40F3F440	4040F5F1
404040F4	F6404040	F4F64040	40F5F240	4040F5F4	404040F5	F0404040	F3F64040	40F3F540	4040F3F5
404040F4	F0404040	F5F14040	40F5F340	4040F3F4	404040F3	F4404040	F4F74040	40F4F540	4040F3F8
404040F3	F4404040	F4F44040	40F4F440	4040F3F6	404040F5	F7404040	F7F14040	40F4F140	4040F4F1
404040F4	F4404040	F5F74040	40F7F340	4040F4F7	404040F4	F1404040	F4F54040	40F8F240	4040F8F2
404040F2	F4404040	F8F14040	40F5F140	4040F6F5	404040F6	F4404040	F7F94040	40F7F440	4040F8F6
404040F6	F5404040	F7F54040	40F8F340	4040F4F8	404040F5	F1404040	F7F24040	40F4F940	4040F4F6
404040F4	F5404040	F4F44040	40F3F540	4040F3F6	404040F5	F3404040	F4F64040	40F4F340	4040F3F9
404040F4	F1404040	F4F04040	40F4F340	4040F4F3	404040F4	F0404040	F3F74040	40F5F540	4040F4F3
404040F4	F3404040	F4F24040	40F4F240	4040F8F4	404040F4	F2404040	F4F14040	40F4F240	4040F4F1
404040F4	F6404040	F4F34040	40F4F540	4040F4F4	404040F3	F6404040	F3F84040	40F4F340	4040F4F1
404040F7	F7404040	F7F64040	40F3F340	4040F4F0	404040F6	F3404040	F4F44040	40F8F140	4040F4F0
404040F4	F0404040	F4F24040	40F4F240	4040F3F6	404040F4	F4404040	F4F24040	40F4F040	4040F2F9
4040F1F1	F5404040	F3F34040	40F3F340	4040F3F5	404040F3	F5404040	F3F44040	40F3F640	4040F3F8
404040F4	F2404040	F3F14040	40F3F640	4040F3F1	404040F6	F6404040	F3F74040	40F7F440	4040F6F1
404040F4	F4404040	F7F24040	40F3F540	4040F3F3	404040F3	F6404040	F3F74040	40F2F540	4040F2F9
404040F7	F4404040	F3F64040	40F3F340	4040F2F5	404040F3	F2404040	F3F44040	40F3F840	4040F3F2
404040F2	F6404040	F6F74040	40F3F540	4040F6F6	404040F6	F7404040	F7F04040	40F6F840	4040F3F1
404040F3	F1404040	F6F74040	40F2F840	4040F3F2	404040F5	F6404040	F3F94040	F1F0F040	4040F2F9
4040F1F0									

RECCED 118 CF FILE 1
LENGTH = 2930 BYTES

40F7F1F8	F240F7F0	F4F740F7	F6F7F640	F7F1F3F0	40F7F0F9	F140F7F0	F4F940F7	F0F7F840	F7F1F8F6
40F7F1F1	F640F7F1	F0F940F7	F6F5F540	F7F1F5F0	40F7F0F1	F440F7F0	F7F140F7	F0F5F440	F7F1F1F5
40F7F1F5	F140F7F0	F5F540F7	F1F6F440	F7F1F3F8	40F7F1F4	F040F7F2	F2F040F7	F1F5F540	F7F2F2F1
40F7F1F9	F140F7F0	F2F740F7	F1F0F140	F7F0F3F5	40F7F0F1	F140F6F5	F9F940F7	F1F3F540	F7F0F5F9
40F7F0F2	F740F7F1	F1F640F7	F1F1F040	F7F1F0F3	40F6F9F7	F540F7F0	F1F340F6	F5F5F640	F7F0F1F8
40F6F9F8	F140F6F9	F4F640F7	F6F3F140	F6F9F2F9	40F6F9F8	F140F7F0	F1F540F5	F5F8F040	F6F9F4F5
40F6F9F5	F040F7F0	F2F640F6	F5F0F540	F6F9F7F3	40F6F8F6	F540F6F9	F6F740F7	F0F1F640	F7F0F4F5
40F7F0F5	F040F7F0	F1F740F6	F5F6F440	F7F0F0F6	40F7F0F0	F240F7F1	F0F340F7	F0F1F040	F6F9F7F9
40F6F9F1	F540F6F9	F5F440F6	F6F6F540	F6F8F9F6	40F6F9F7	F640F6F8	F4F540F6	F7F9F440	F6F8F0F8
40F6F8F9	F340F6F9	F3F440F6	F6F3F540	F6F9F8F6	40F6F8F4	F240F6F8	F4F540F6	F8F7F840	F6F8F9F8
40F6F8F6	F740F6F6	F2F440F6	F5F3F340	F6F8F7F0	40F6F8F6	F740F6F8	F7F240F6	F5F5F540	F6F9F2F9
40F6F9F5	F740F6F9	F3F340F6	F5F2F640	F6F9F2F1	40F6F9F0	F340F6F9	F5F240F6	F5F4F740	F6F8F5F2
40F6F9F2	F040F7F0	F5F540F6	F6F5F440	F7F0F1F3	40F6F9F5	F040F6F9	F9F340F6	F5F2F540	F6F5F4F3
40F6F9F7	F640F6F9	F5F640F6	F5F5F540	F7F0F5F2	40F6F8F7	F440F6F9	F3F640F6	F6F6F540	F6F7F4F7
40F6F9F2	F340F6F8	F2F240F6	F6F6F840	F6F8F5F4	40F6F8F5	F240F6F8	F6F640F6	F8F2F940	F6F8F3F0
40F6F7F4	F540F6F8	F0F640F6	F6F0F240	F6F8F2F2	40F6F8F5	F540F6F8	F9F940F6	F8F0F140	F6F8F0F7
40F6F8F2	F540F6F6	F6F540F6	F6F1F340	F6F8F5F0	40F6F7F4	F540F6F7	F6F740F6	F6F4F440	F6F8F5F2
40F6F7F8	F540F6F7	F7F140F6	F7F1F040	F6F8F2F9	40F6F6F5	F540F6F6	F3F640F6	F6F1F640	F6F6F6F7
40F6F5F5	F440F6F6	F4F640F6	F6F2F140	F6F6F1F7	40F6F6F7	F740F6F5	F8F840F6	F6F4F440	F6F6F5F4
40F6F6F1	F640F6F5	F5F440F6	F6F3F040	F6F6F1F8	40F6F6F5	F540F6F6	F3F840F6	F6F7F440	F6F7F0F8
40F6F7F3	F640F6F6	F5F440F6	F7F1F540	F6F7F0F3	40F6F6F8	F540F6F7	F2F540F6	F7F7F040	F6F7F4F8
40F6F8F3	F440F6F7	F7F740F6	F7F6F640	F6F7F5F5	40F6F8F1	F640F6F8	F2F540F6	F7F7F540	F6F7F5F2
40F6F7F4	F140F6F7	F7F640F6	F6F6F440	F6F7F8F5	40F6F8F2	F240F6F8	F5F440F6	F8F6F240	F6F9F8F0
40F6F8F9	F540F6F8	F6F440F6	F5F2F340	F6F8F8F1	40F6F9F2	F040F6F9	F5F140F6	F5F5F040	F7F0F5F6
40F6F9F6	F540F7F0	F6F740F6	F5F6F640	F7F0F3F6	40F6F9F5	F540F6F9	F8F240F6	F5F2F640	F6F9F7F3
40F6F9F4	F640F7F0	F2F140F6	F5F8F440	F7F0F0F7	40F7F0F0	F740F7F1	F9F740F7	F1F0F940	F6F8F7F6
40F6F8F9	F740F6F8	F4F540F6	F6F5F540	F6F8F4F6	40F6F9F2	F640F6F5	F9F940F6	F5F6F640	F7F0F2F8
40F7F0F3	F640F7F0	F7F440F7	F1F6F340	F7F0F5F5	40F7F0F9	F540F7F1	F2F140F7	F0F5F240	F7F1F8F9
40F7F1F0	F540F7F0	F0F140F7	F6F7F640	F7F0F5F4	40F7F1F0	F640F7F0	F0F540F7	F0F3F340	F7F0F2F4
40F7F0F2	F440F6F9	F0F940F6	F5F5F640	F6F8F9F5	40F6F9F4	F640F6F9	F8F140F6	F5F1F640	F7F0F4F2
40F6F2F0	F5F1F0F9	F4F6F1F2	F2F4F6F4	F1F2F3F7	F0F1F2F0	F9F4F1F0	F9F4F1F0	F5F6F9F1	F0F1F5F3
40F6F5F1	F640F7F3	F6F640F7	F1F4F640	F7F0F5F9	40F6F9F5	F240F7F1	F9F540F7	F1F4F740	F7F1F4F0
40F7F0F5	F740F7F2	F3F640F7	F3F4F440	F7F1F5F8	40F7F3F1	F540F7F3	F2F240F7	F0F9F340	F7F1F7F4
40F7F1F0	F140F7F0	F3F140F7	F6F2F640	F7F0F2F6	40F7F1F2	F540F7F1	F1F840F7	F0F6F740	F7F0F5F1
40F7F1F8	F140F7F1	F7F540F7	F1F4F240	F7F1F8F6	40F7F1F7	F540F7F1	F6F440F6	F5F5F640	F7F1F4F0
40F7F2F4	F440F7F0	F5F640F7	F1F6F140	F7F2F6F7	40F7F1F9	F140F7F1	F0F640F7	F1F5F340	F7F2F5F0
40F7F1F7	F240F7F2	F4F040F7	F1F6F540	F7F3F9F7	40F6F5F3	F1404040	F9F94040	40F3F140	4040F6F9
404040F2	F5404040	F3F44040	40F5F640	4040F3F4	404040F6	F1404040	F3F14040	40F3F140	4040F6F6
404040F7	F2404040	F3F44040	40F2F740	4040F2F6	404040F3	F1404040	F7F14040	40F2F540	4040F4F0
404040F3	F7404040	F3F34040	40F3F140	4040F3F5	404040F4	F1404040	F2F94040	40F3F040	4040F2F4
404040F3	F2404040	F3F74040	40F2F540	4040F3F6	404040F3	F1404040	F7F64040	40F6F240	4040F3F7
404040F3	F4404040	F3F54040	40F3F440	4040F2F6	404040F3	F6404040	F4F24040	40F7F540	4040F3F8
404040F2	F6404040	F3F54040	40F6F540	4040F7F0	404040F4	F0404040	F3F54040	40F3F040	4040F3F7
404040F3	F2404040	F3F04040	40F4F640	4040F4F1	404040F7	F6404040	F4F04040	40F3F740	4040F3F6

F5F0F6F4	F1F5F0F6	F4F0F5F0	F4F3F3F5	F0F6F3F6	F5F0F6F3	F4F5F0F6	F3F2F5F0	F4F3F0F5	F0F6F2F7
F5F0F6F2	F5F5F0F6	F2F2F5F0	F4F1F5F5	F0F6F1F4	F5F0F6F0	F5F5F0F6	F0F3F3F0	F5F9F8F5	F0F5F9F3
F5F0F5F8	F5F5F0F5	F8F2F5F0	F5F7F7F5	F0F5F7F1	F5F0F5F6	F4F5F0F5	F6F0F5F0	F5F5F4F5	F0F5F4F9
F5F0F5F4	F3F5F0F5	F3F7F5F0	F5F3F1F5	F0F5F2F5	F5F0F5F1	F5F5F0F5	F1F3F5F0	F5F0F7F5	F0F5F0F1
F5F0F4F9	F4F5F0F4	F2F8F5F0	F4F8F2F5	F0F4F7F5	F5F0F4F6	F5F5F0F4	F6F2F5F0	F4F5F6F5	F0F4F4F9
F5F0F4F4	F2F5F0F4	F3F6F5F0	F4F2F5F5	F0F4F2F2	F5F0F4F1	F5F5F0F4	F0F5F5F0	F4F0F2F5	F0F3F5F5
F5F0F3F8	F7F5F0F3	F7F5F5F0	F3F7F1F5	F0F3F6F3	F5F0F3F5	F5F5F0F3	F4F6F5F0	F3F3F5F5	F0F3F3F0
F5F0F3F2	F2F5F0F3	F1F3F5F0	F2F0F5F5	F0F2F9F7	F5F0F2F8	F5F5F0F2	F8F1F5F0	F2F7F3F5	F0F2F6F5
F5F0F2F5	F7F5F0F2	F5F0F5F0	F2F4F2F5	F0F2F3F4	F5F0F2F2	F7F5F0F2	F1F5F5F0	F2F1F1F5	F0F2F0F4
F5F0F1F5	F7F5F0F1	F8F5F5F0	F1F4F2F5	F0F1F7F5	F5F0F1F6	F7F5F0F1	F6F0F5F0	F1F5F3F5	F0F1F4F6
F5F0F1F3	F5F5F0F1	F3F2F5F0	F1F2F5F5	F0F1F1F5	F5F0F1F1	F3F5F0F1	F0F6F5F0	F1F0F0F5	F0F0F5F3
F5F0F0F8	F7F5F0F0	F6F1F5F0	F0F7F5F5	F0F0F6F5	F5F0F0F6	F3F5F0F0	F5F7F5F0	F0F5F1F5	F0F0F4F6
F5F0F0F4	F0F5F0F0	F3F5F5F0	F6F2F5F5	F0F0F2F4	F5F0F0F1	F5F5F0F0	F1F4F5F0	F0F1F0F5	F0F0F0F6
F5F0F0F0	F2F4F9F5	F5F7F4F5	F5F5F3F4	F5F9F8F5	F4F9F9F8	F4F4F5F0	F7F5F4F9	F5F7F4F4	F5F5F5F5
F4F5F5F6	F4F4F9F5	F5F8F4F5	F5F5F3F4	F5F9F4F8	F4F9F9F4	F2F4F9F5	F3F7F4F5	F5F3F2F4	F5F9F2F5
F4F5F5F2	F3F4F9F5	F1F9F4F5	F5F1F0F4	F5F9F1F4	F4F5F5F1	F2F4F9F5	F1F2F4F5	F5F1F3F4	F5F9F1F5
F4F5F5F1	F2F4F9F5	F0F5F4F5	F5F0F7F4	F5F9F0F5	F4F9F5F0	F3F4F9F5	F0F1F4F5	F5F5F5F4	F5F8F5F8
F4F5F5F5	F6F4F9F5	F5F5F4F5	F5F5F4F5	F5F8F9F3	F4F9F8F5	F2F4F9F5	F9F2F4F5	F5F5F1F4	F5F8F5F1
F4F5F5F5	F3F4F9F5	F5F9F5F0	F0F0F5F5	F0F0F1F1	F5F0F0F1	F5F5F0F0	F2F5F5F0	F0F3F1F5	F0F0F3F9
F5F0F0F4	F4F5F0F0	F5F3F5F0	F0F6F1F5	F0F0F6F6	F5F0F0F7	F5F5F0F0	F8F4F5F0	F0F9F3F5	F0F1F0F1
F5F0F1F0	F5F5F0F1	F1F8F5F0	F1F2F7F5	F0F1F3F6	F5F0F1F4	F5F5F0F1	F5F5F5F0	F1F6F5F5	F0F1F7F4
F5F0F1F8	F4F5F0F1	F5F4F5F0	F2F0F4F5	F0F2F1F4	F5F0F2F2	F4F5F0F2	F3F4F5F0	F2F4F4F5	F0F2F5F4
F5F0F2F6	F3F5F0F2	F7F1F5F0	F2F8F0F5	F0F2F8F5	F5F0F2F5	F5F5F0F3	F0F7F5F0	F3F1F7F5	F0F3F2F6
F5F0F3F3	F5F5F0F3	F4F4F5F0	F2F5F3F5	F0F3F6F2	F5F0F3F7	F1F5F0F3	F8F1F5F0	F3F8F9F5	F0F3F5F9
F5F0F4F0	F7F5F0F4	F1F7F5F0	F4F2F5F5	F0F4F3F4	F5F0F4F4	F3F5F0F4	F5F2F5F0	F4F6F1F5	F0F4F7F0
F5F0F4F7	F5F5F0F4	F6F8F5F0	F4F5F7F5	F0F5F0F7	F5F0F5F1	F5F5F0F5	F2F5F5F0	F5F3F4F5	F0F5F4F3
F5F0F5F5	F3F5F0F5	F4F2F5F0	F5F7F1F5	F0F5F8F1	F5F0F5F5	F0F5F0F5	F9F8F5F0	F4F0F5F5	F0F6F1F2
F5F0F6F1	F5F5F0F6	F2F5F5F0	F4F3F2F5	F0F6F3F5	F5F0F6F4	F4F5F0F6	F5F3F5F0	F6F5F0F5	F0F6F2F7
F5F0F6F7	F4F5F0F6	F8F1F5F0	F4F5F7F5	F0F6F5F4	F5F0F7F0	F1F5F0F7	F0F8F5F0	F7F1F4F5	F0F7F2F1
F5F0F7F2	F7F5F0F7	F3F4F5F0	F1F4F0F5	F0F7F4F6	F5F0F7F5	F3F5F0F7	F5F5F5F0	F7F6F5F5	F0F7F7F1
F5F0F7F7	F7F5F0F7	F8F24040	4040CF140	404040F1	40404040	F1404040	40F14040	4040F140	404040F1
40404040	F1404040	40F14040	4040CF140	404040F1	40404040	F1404040	40F14040	4040F140	404040F1
40404040	F1404040	40F14040	4040CF140	404040F1	40404040	F1404040	40F14040	4040F1	

[illegible]

RECORD = 118 OF FILE 140
LENGTH = 2930 BYTES

118 RECCFES IN FILE 140 CF TAPE